

## CO<sub>2</sub> Transoral Laser Microsurgery (TLM) Reference Article List

### Lumenis Global Sales Offices

#### America

##### Santa Clara, CA

Tel + 1 408 764 3000  
+ 1 877 586 3647  
Fax + 1 408 764 3999

#### Europe, Middle East & Africa

##### Dreieich-Dreieichenhain, Germany

Tel + 49 6103 8335 0  
Fax + 49 6103 8335 300

##### Formello (RM), Italy

Tel + 39 06 90 75 230  
Fax + 39 06 90 75 269

##### London, UK

Tel + 44 20 8324 4200  
Fax + 44 20 8324 4222

#### Asia-Pacific

##### Beijing, China

Tel + 86 10 6510 2620  
Fax + 86 10 6510 2621

##### Hong Kong

Tel + 852 2174 2800  
Fax + 852 2722 5151

##### Gurgaon, India

Tel + 91 124 422 33 80  
Fax + 91 124 422 33 83

#### Japan

##### Tokyo, Japan

Tel + 81 3 5789 8300  
Fax + 81 3 5789 8301

Internet: [www.lumenis.com](http://www.lumenis.com)

Email: [information@lumenis.com](mailto:information@lumenis.com)

Holsinger FC, Weber R. Swing of the surgical pendulum: A return to surgery for treatment of head and neck cancer in the 21st century?: Int. J. Radiation Oncology Biol. Phys., Vol. 69, 2007; No. 2, Supplement, S129 – S131

Hinni ML, Salassa JR, Haughey B, Steiner W, et. al. Transoral Laser Microsurgery for Advanced Laryngeal Cancer. Arch Otolaryngol Head Neck Surgery, Dec 2007; Vol. 133 (No.12)

Sjögren E, et. al. Voice Outcome in T1a Midcord Glottic Carcinoma: Laser Surgery vs. Radiation; Arch Otolaryngol Head Neck Surgery, Sep 2008; Vol. 134 (No.9)

Schrijvers ML, et. al. Higher Laryngeal Preservation Rate After CO<sub>2</sub> Laser Surgery Compared With Radiotherapy in T1A Glottic Laryngeal Carcinoma. Head & Neck, June 2009; DOI 10.1002/hed

Steiner W, Ambrosch P. Endoscopic laser surgery of the upper aerodigestive tract. 2000 Thieme New York, ISBN 0-86577-996-1

Rich JT, et. al. Transoral Laser Microsurgery (TLM) +/- adjuvant therapy for advanced stage oropharyngeal cancer: outcome and prognostic factors. Laryngoscope, July 2009

Preuss SF, Cramer K, Klussmann JP, Eckel H, Guntinas-Lichius O. Transoral laser surgery for laryngeal cancer: Outcome, complications and prognostic factors in 275 patients. European Journal Surgery Oncology, 2008

Roh JL, Kim DH, Park CI. Voice, Swallowing and Quality of Life in Patients After Transoral Laser Surgery for Supraglottic Carcinoma. Journal of Surgical Oncology, 2008 Wiley-Liss

Cohen SM, Garrett CG, Dupont WD, Ossoff RH, Courey MS. Voice-related quality of life in T1 glottic cancer: irradiation versus endoscopic excision. Ann Otol Rhinol Laryngol. 2006; 115 (8): 581-586

Remacle M, Hassan F, Cohen D, Lawson G, Delos M, 2005: New computer-guided scanner for improving CO<sub>2</sub> laser-assisted microincision. European Arch Otorhinolaryngol, 2005; 262: 113-119

Remacle M, Lawson G, Nollevaux MC, Delos M. Current state of scanning micromanipulator applications with the carbon dioxide laser. Annals of Otolaryngology, Rhinology & Laryngology, 2008; 117(4):239-244

Roediger FC, Orloff LA, Courey MS. Adult subglottic stenosis: management with laser incisions and Mitomycin-C. Laryngoscope, Sept 2008; 118

Brandenburg JH. Laser cordotomy versus radiotherapy: an objective cost analysis. Ann Otol Rhinol Laryngol 2001; 110: 312-318

Goor KM, Peeters AJ, Mahieu HF, et. al. Cordectomy by CO<sub>2</sub> laser or radiotherapy for small T1a glottic carcinomas: cost, local control, survival, quality of life, and voice quality: Head Neck 2007; 29:128-136

Cragle SP, Brandenburg JH. Laser cordectomy or radiotherapy: cure rates, communication, and cost. Otolaryngol Head Neck Surgery. 1993; 108(6):648-654